

Tear film measurements in four different ethnic groups: Malay, Chinese, Indian and Nigerian

Madhavendra Bhandari¹, Boo Sheau Yun²

¹School of Optometry, Faculty of Medicine and Health Sciences, UCSI University, Kuala Lumpur 52200, Malaysia

²Twintech International University Collage of Technology, Kuala Lumpur 52200, Malaysia

Correspondence to: Madhavendra Bhandari. School of Optometry, Faculty of Medicine and Health Sciences, UCSI University, Kuala Lumpur 52200, Malaysia. madhavendraopto@yahoo.co.in

Received: 2019-11-18 Accepted: 2021-01-13

Abstract

• **AIM:** To compare the non-invasive tear film break-up time (NIBUT), tear break up time (TBUT), basal tear secretion (BTS) and blink rate in four ethnic groups: Malay, Chinese, Indian and Nigerian.

• **METHODS:** Totally 120 healthy (61 males and 59 females) subjects (without dry eye symptoms and ocular surface disorder) with the age 20 to 39 years were recruited; 30 were Malays, 30 were Chinese, 31 were Indians and 29 were Nigerians. Based on McMonnies questionnaire and clinical examination, normal subjects were selected. NIBUT, TBUT, BTS were assessed in only one eye (right) of each subject and blink rate was also assessed.

• **RESULTS:** There was significant difference in the NIBUT, TBUT, BTS and blink rate among 4 different ethnic groups ($P=0.018$, 0.001 , 0.011 , and 0.004 respectively). No statistically significant difference of NIBUT, TBUT, BTS and blink rate was found between the genders among different ethnic groups. Indian had higher median for NIBUT (10 ± 6 s), TBUT (7 ± 5 s) and BTS (20 ± 20 mm) than other races. Chinese had lower median for NIBUT (7.5 ± 4 s) and TBUT (4 ± 2 s) while Malay had for BTS (9.5 ± 16 mm) among the groups. There was no significant correlation of blink rate with NIBUT ($r=-0.119$, $P=0.195$), TBUT ($r=-0.086$, $P=0.352$), and BTS ($r=-0.123$, $P=0.180$) respectively.

• **CONCLUSION:** The tear-film measurement values are variability in four ethnic groups.

• **KEYWORDS:** non-invasive tear film break-up time; tear break up time; basic tear secretion; blink rate; ethnicity

DOI:10.18240/ier.2021.02.02

Citation: Bhandari M, Sheau YB. Tear film measurements in four different ethnic groups: Malay, Chinese, Indian and Nigerian. *Int Eye Res* 2021;2(2):69-73

INTRODUCTION

Measurements of the stability and volume of tear film are important while assessing the ocular surface in cases of dry eye syndrome. Dry eye syndrome and tear film dysfunction are synonymous. Tear film dysfunction leads to ocular complications such as excessive tearing, conjunctival and corneal surface damage, and vision changes^[1-2]. Therefore, the tear stability and volume is essential in assessing contact lens fitting, cataract and refractive surgery, and also to know the outcome of dry eye treatment^[3-9]. The normal values of tear stability and volume may differ in different ethnic groups. Differences in eyelid anatomy (e.g. palpebral aperture size and lid margin apposition against the ocular surface) between Asians and non-Asians may contribute to variability in tear film thickness and tear-film stability^[10]. Studies suggesting wider palpebral aperture sizes and palpebral aperture width lead to more exposure and greater disruption in tear film^[11-12].

There are separate reports on tear stability and tear volume in different ethnic groups living in different environments^[13-20]. However, due to the difference in instrument designs and observational procedures; there are differences in tear film measurements. The purpose of this study was to determine the tear stability and tear volume in Chinese, Malay, Indian and Nigerian ethnic groups living in the same climate.

SUBJECTS AND METHODS

Ethical Approval The study was carried out in the Optometry Clinic at Twintech International University College of Technology, Faculty of Optometry, Kuala Lumpur, Malaysia. Informed consent was obtained from all participants prior to any measurement. The study was conducted according to the tenets of Declaration of Helsinki.

This was a prospective cross sectional study. A total of 120 subjects (age range 20-39 years) were recruited. The subjects were from 4 different ethnicities; 30 Malays, 30 Chinese, 31 Indians and 29 Nigerians. All the subjects were students, 3 ethnics groups; Malays, Chinese, and Indians were Malaysian whereas, Nigerians were from Nigeria, studying in Malaysia under same university.

The inclusion criteria set for this study was as follow: age between 20 to 39 years, currently not on any medication, not contact lens wearers, have no history of any ocular surgery, have habitual visual acuity of 20/30 or better, have normal appearance of all corneal layers on slit lamp examination and with no corneal lesions. Individuals with autoimmune disorders, and females during pregnancy, menstruation, lactation, and hormone replacement therapy were excluded. Subjects having a history of dry eye and Mc-Monnies questionnaire score more than 14.5 were also excluded from this study to rule out dry eye^[21].

Habitual visual acuity was measured using a Snellen chart. Tear film status was then evaluated. Blink rate was measured first. The number of blinks made per minute was recorded. Non-invasive tear film break-up time (NIBUT), tear break up time (TBUT), basic tear secretion (BTS) were assessed in only one eye (right) of each subject.

The NIBUT test, which measures tear stability without the introduction of a foreign substance into the tear film^[22], was assessed next by using a Shin-Nippon S0-21 keratometer. The subject was seated carefully and the keratometer well adjusted for the right eye with the mires in focus. The subject was asked to blink once and then refrain from blinking. A stopwatch was started after the last complete blink. At the first sign of any distortion of the mires, the stopwatch was stopped and the time noted. If the subject blinked between measurements, the test was repeated after several blinks. The time interval between the last blink and the first sign of mire distortion was recorded in seconds as the NIBUT. The test was repeated three times to increase the reliability and the average was then taken^[14].

TBUT is the time needed for the appearance of dark spots of the tear film after blinking of the fluorescein treated eye^[22]. For the measurement of TBUT, the upper eyelid of the right eye was slightly lifted, and the fluorescein strip was then used to stain the tear film. The subject was asked to blink a few times for homogenous mixing of fluorescein in tear film. Then, the tear film was observed under cobalt-blue filter. The stopwatch was started after the last blink and the appearance of dry spots on the corneal surface was detected and recorded in seconds. An average of three readings was taken.

After resting for 10min, BTS was measured using Schirmers strips (Whatman no 41 filter paper, tear touch). A drop of tropical anaesthesia (0.5%, Alcaine) was instilled in the tested eye. After 2min, the Schirmers strip was placed at the lateral one third of the lower eyelid of the eye. The subject was asked to blink normally. After 5min, the strip was removed. The length of the wetted strip from the fold was measured.

All the subjects were tested between 10 a.m.-12 p.m., at least after 2h of waking. Tear film measurements were performed in the same room to maintain a relatively constant temperature




Figure 1 Median of blink rate, NIBUT, TBUT and BTS in ethnic groups.

and humidity. Slit lamp biomicroscopy of the anterior segment (lids, cornea, and conjunctiva) was also performed. Meibomian gland dysfunction was recorded according to Efron grading scale. The grade 0-1 was considered as normal in this study^[23].

Statistical Analysis Statistical analyses were done by using SPSS version 16.0 statistical software and MS Excel 2007. Test of normality was performed by Kolmogorov-Smirnov test. The measurement variables were not normally distributed. Kruskal-Wallis test was performed among four different ethnic groups. Mann-Whitney test was performed between the ethnic groups and sex groups. To evaluate correlation between variables, Spearman correlation coefficient was used. A $P \leq 0.05$ was set for statistical significance.

RESULTS

A total of 120 subjects (mean: 23.07 ± 2.69 y; range 20-39y) were examined. There were 61 (50.8%) male and 59 (49.2%) female subjects. Thirty (25%) subjects were Malays, 30 (25%) were Chinese, 31 (25.8%) were Indians and 29 (24.2%) were Nigerians. There was no statistically significant difference in age among ethnic groups. There was statistically significant difference in the NIBUT ($P=0.018$), TBUT ($P=0.001$), BTS ($P=0.011$) and blink rate ($P=0.004$) among ethnic groups (Table 1, Figure 1).

Indians had the highest NIBUT, TBUT, and BTS whereas Chinese had the lowest mean NIBUT and TBUT (Table 1). The NIBUT, TBUT and BTS amongst Indians were significantly higher compared to Malays, and Chinese (Table 2). The blink rate for Malays was significantly higher than for Indians, Chinese and Nigerians (Table 2).

There was no statistically significant difference in age ($P=0.095$), blink rate ($P=0.881$), NIBUT ($P=0.114$), TBUT ($P=0.841$), and BTS ($P=0.348$) between genders (Table 3). Furthermore, there was no significant difference of tear film measurements (blink rate, NIBUT, TBUT, and basic tear secretion) with gender in Malay, Chinese, Indian, and Nigerian. Table 4 shows correlation between tear film variables, and age. The degree of association between NIBUT, TBUT and BTS

Table 1 Comparison of age and tear film measurements among ethnic groups median±IQR

Variables	Total (n=120)	Malay (n=30)	Chinese (n=30)	Indian (n=31)	Nigerian (n=29)	P
Age	23±3	23±2	23±1	23±2	22±4	0.093
Blink rate (blinks/min)	12±6	16±8	12±6	12±6	12±4	0.004
NIBUT (s)	8±4	8±4	7.5±4	10±6	8±4	0.018
TBUT (s)	5.3±4	5±3	4±2	7±5	6±4	0.001
Basic tear secretion (mm)	13±14	9.5±16	11.5±10	20±20	13±12	0.011

Table 2 Comparison (P) of tear film measurements between ethnic groups

Groups	Variables	Chinese	Indian	Nigerian
Malay	Blink rate	0.02	0.03	0.001
	NIBUT	0.384	0.046	0.531
	TBUT	0.184	0.01	0.173
	Basic tear secretion	0.407	0.003	0.129
Indian	Blink rate	0.791	-	-
	NIBUT	0.002	-	-
	TBUT	0.001	-	-
	Basic tear secretion	0.01	-	-
Nigerian	Blink rate	0.214	0.106	-
	NIBUT	0.086	0.112	-
	TBUT	0.004	0.101	-
	Basic tear secretion	0.417	0.079	-

NIBUT: Non-invasive tear film break-up time; TBUT: Tear break up time.

Table 3 Comparison of age and tear film measurements between genders

Parameters	Genders	Median±IQR	P
Age (y)	M (n=61)	23±2	0.095
	F (n=59)	23±2	
Blink rate (blinks/min)	M (n=61)	12±7	0.881
	F (n=59)	12±6	
NIBUT (s)	M (n=61)	9±4	0.114
	F (n=59)	8±5	
TBUT (s)	M (n=61)	5±4	0.841
	F (n=59)	6±4	
Basic tear secretion (mm)	M (n=61)	13±16	0.348
	F (n=59)	12±14	

was significant. There was, however, no significant correlation of blink rate with NIBUT ($r=-0.119$, $P=0.195$), TBUT ($r=-0.086$, $P=0.352$) and BTS ($r=-0.123$, $P=0.180$) respectively.

DISCUSSION

Our results shown that Indian had highest tear film measurements (NIBUT: 10s; TBUT: 7s and BTS: 20 mm) among studied groups. While Chinese had lowest median for NIBUT (7.5s), TBUT (4s) compared to the others ethnic. Furthermore, Malay had lowest median for BTS (9.5 mm) and highest for blink rate (16 blinks/min). Between the genders, there was no statistically significant difference of tear film measurement values in Malay, Chinese, Indian and Nigerian.

Based on previous studies, humidity and temperature has been shown to affect the tear film stability^[24-27]. Therefore, in the present study, all measurements were performed in same room to maintain a relatively constant temperature and humidity. Earlier studies have shown that there is a diurnal variation of the tear film^[28-30]. Shen *et al*^[28] have reported that the tear meniscus increased at eye opening, after overnight sleep, compared with before sleep. They also reported that the stability of the pre-corneal tear film varied over a single 12-hours period. Another study suggested that the tear stability is low early in the morning, and tends to rise to equilibrium level between 10:00 a.m. and 12 p.m.^[29]. However, Srinivasan and colleagues showed no significant diurnal changes in the bare-eye tear film surface quality^[30]. In order to avoid diurnal variations, the present study was performed between 10 a.m.-12 p.m. There is an influence of corneal hydration due to reduce in tear film evaporation within 2 hours of waking. Thus, tear film procedures were performed in present study after 2h of waking^[31].

The present study showed that different ethnic groups had different tear stability. This is similar to some previous studies which show that there is variability in TBUT values in different ethnic groups^[15,17]. The NIBUT, TBUT, and BTS of Indian ethnic group were significantly higher than Chinese, Malay, and Nigerian in the present study. Patel *et al*^[15] also reported similar findings in that the mean NIBUT of Indians was higher compared to Chinese and African while Chinese had lower value of NIBUT. Cho & Brown noted that the mean TBUT of Hong Kong-Chinese was approximately 8s and about 70% of Hong Kong-Chinese had TBUT's less than 10s^[20]. Among Malaysian, present study showed Chinese had lower TBUT and NIBUT, also similar finding by Amaechi *et al*^[18] and Sharanjeet *et al*^[19] suggesting Chinese had lower TBUT and NIBUT respectively. Studies reported morphological difference of periorbital features like palpebral aperture size and width in different ethnic groups^[32-34]. Thus, it influences the tear film measurement^[11-12]. Packiriswamy *et al*. reported the mean palpebral fissure height values in Malaysian both male (Indian 10.9 mm; Malay, 10 mm; Chinese 9.6 mm) and female (Indian 11.3mm; Malay 1.70 mm; Chinese 10.3 mm)^[32]. Interestingly, present study as well as previous studies suggested tear film

Table 4 Correlation between study variables

Variables	Blink rate	NIBUT	TBUT	BTS
Age	$r=-0.931, P=0.311$	$r=-0.015, P=0.872$	$r=-0.32, P=0.729$	$r=-0.54, P=0.556$
Blink rate	-	$r=-0.119, P=0.195$	$r=-0.086, P=0.352$	$r=-0.123, P=0.180$
TBUT	-	$r=0.786, P=0.001$	-	$r=0.384, P=0.001$
NIBUT	-	-	-	$r=0.345, P=0.001$

NIBUT: Non-invasive tear film break-up time; TBUT: Tear break up time; BTS: Basic tear secretion.

Table 5 Comparison of present study with previous studies based on ethnic groups

Studies	Malays		Chinese		Indian		Nigerian	
	TBUT (s)	NIBUT (s)	TBUT (s)	NIBUT (s)	TBUT (s)	NIBUT (s)	TBUT (s)	NIBUT (s)
Madugil <i>et al</i> ^[13]					13.87±4.76			
Cho ^[14]	-	-	-	16±9.4	-	-	-	-
Patel <i>et al</i> ^[15]	-	-	-	9.8±3.90	-	16.4±6.9	-	11.8±5.9
Mohidin <i>et al</i> ^[16]	-	15.8±9.4	-	-	-	-	-	-
Jamaliah <i>et al</i> ^[17]	9.65±4.14	-	8.61±3.73	-	8.92±2.92	-	-	-
Amaechi <i>et al</i> ^[18]	-	-	-	-	-	-	15.2±3.1	15.3±3.0
Sharanjeet <i>et al</i> ^[19]		7.74±3.34		7.15±3.38		7.24±2.67		
Present study	5±3	8±4	4±2	7.5±4	7±5	10±6	6±4	8±4

stability increases with greater palpebral fissure dimension while comparing among the Malaysian ethnics groups^[18-19,32-34]. This present study reported no significant difference in blink rate, NIBUT, TBUT and BTS between males and females. This is also supported by earlier studies with similar results between the genders^[15-16,18,20]. However, few reports showed the majority of dry-eye sufferers are women, and female gender has been identified as a risk factor for dry-eye development^[35-36]. The difference in sex hormones that influences the anatomical and functional structures of the ocular surface, and tear film components^[36-37].

Blink rate was not correlated with NIBUT, TBUT and BTS in this study. There are a number of factors that can alter blink rate and have a profound impact on the ocular surface. Multiple factors can contribute to changes in blink rate, for example, thickness of the lipid layer, rate of eye drying, humidity, temperature, pollutants and visual tasks^[38-41]. Thus, although blink rate is fairly constant for any individual, it can be affected by external conditions.

Although, the results from the present study indicate that there are reduced values of TBUT and NIBUT compared to previous studies, the results are comparable within the ethnic groups (Table 5). Interestingly, there is wide variability in normal values of TBUT and NIBUT between previous studies and recent studies. The recent studies have reported that the values for TBUT and NIBUT were lesser compared to those from earlier studies. This reduction of values of TBUT and NIBUT in recent studies may be mainly due to environmental factors and other factors like different age population distribution, and methodology.

There are few limitations of this study. First, there was no record of temperature and humidity, even though all the examinations were performed in same room to control the changes in tear. Second, the differences in instrument designs and observation procedures could cause the discrepancies in measurement values of tear film while comparing with previous studies. A modified perimeter bowl instrument was used by Mohidin *et al*^[16] and Cho to measure the NIBUT^[16]. Patel *et al*^[15] and Amaechi *et al*^[18] had measured NIBUT by keratometer whereas Sharanjeet-Kaur *et al*^[19] used tearscope plus. The study showed that tear film measurements are different in the 4 different ethnic groups (Malay, Chinese, Indian and Nigerian). Comparatively, Indians has highest tear film measurement values compared to other ethnic groups studied. The Chinese had lowest NIBUT and TBUT compared with the other ethnic groups. Therefore, the norm values for tear stability and tear volume should be established based on the ethnic distribution.

ACKNOWLEDGEMENTS

Conflicts of Interest: Bhandari M, None; Sheau YB, None.

Peer Review File Available at: http://ier.ijo.cn/gjykier/ch/reader/download_attache_file.aspx?seq_id=20210625162023001&flag=1&journal_id=gjykier&year_id=2021&issue=2

REFERENCES

1 Johnson ME, Murphy PJ. Changes in the tear film and ocular surface from dry eye syndrome. *Prog Retin Eye Res* 2004;23(4):449-474.
2 Koh S, Tung CI, Inoue Y, Jhanji V. Effects of tear film dynamics on quality of vision. *Br J Ophthalmol* 2018;102(12):1615-1620.

- 3 Ridder WH, Tomlinson A, Huang JF, Li J. Impaired visual performance in patients with dry eye. *Ocul Surf* 2011;9(1):42-55.
- 4 Grubbs JR, Tolleson-Rinehart S, Huynh K, Davis RM. A review of quality of life measures in dry eye questionnaires. *Cornea* 2014;33(2):215-218.
- 5 Cohen E, Spierer O. Dry eye post-laser-assisted *in situ* keratomileusis: major review and latest updates. *J Ophthalmol* 2018;2018:4903831.
- 6 Iglesias E, Sajjani R, Levitt RC, Sarantopoulos CD, Galor A. Epidemiology of persistent dry eye-like symptoms after cataract surgery. *Cornea* 2018;37(7):893-898.
- 7 Oh T, Jung Y, Chang D, Kim J, Kim H. Changes in the tear film and ocular surface after cataract surgery. *Jpn J Ophthalmol* 2012;56(2):113-118.
- 8 Siddireddy JS, Vijay AK, Tan J, Willcox M. The eyelids and tear film in contact lens discomfort. *Cont Lens Anterior Eye* 2018;41(2):144-153.
- 9 Roberts CW, Camiglia PE, Brazzo BG. Comparison of topical cyclosporine, punctal occlusion, and a combination for the treatment of dry eye. *Cornea* 2007;26(7):805-809.
- 10 Blake CR, Lai WW, Edward DP. Racial and ethnic differences in ocular anatomy. *Int Ophthalmol Clin* 2003;43(4):9-25.
- 11 Lam CSY, Loran DFC. Designing contact lenses for oriental eyes. *J Br Contact Lens Assoc* 1991;14(3):109-114.
- 12 Brasil MV, Brasil OF, Vieira RP, Vaisman M, Amaral Filho OM. Tear film analysis and its relation with palpebral fissure height and exophthalmos in Graves' ophthalmopathy. *Arq Bras Oftalmol* 2005;68(5):615-618.
- 13 Maudgil SS, Khurana AK, Singh M, Parmar IP, Ahluwalia BK. Tear film flow and stability in normal Indian subjects. *Indian J Ophthalmol* 1989;37(4):182-183.
- 14 Cho P. Reliability of a portable noninvasive tear break-up time test on Hong Kong-Chinese. *Optom Vis Sci* 1993;70(12):1049-1054.
- 15 Patel S, Virhia SK, Farrell P. Stability of the precorneal tear film in Chinese, African, Indian, and Caucasian eyes. *Optom Vis Sci* 1995;72(12):911-915.
- 16 Mohidin N, Bay TC, Yap M. Non-invasive tear break-up time in normal Malays. *Clin Exp Optom* 2002;85(1):37-41.
- 17 Jamaliah R, Fathilah J. Prevalence of dry eye in university *Malaya* medical centre. *Med J Malays* 2002;57(4):390-397.
- 18 Amaechi OU, Osunwoke CM. The relation between invasive and non-invasive tear break up time in young adult. *JNOA* 2004;11:29-32.
- 19 Sharanjeet-Kaur, Ho CY, Mutalib HA, Ghazali AR. The relationship between tear ferning patterns and non-invasive tear break-up time in normal Asian population. *J Optom* 2016;9(3):175-181.
- 20 Cho P, Brown B. Review of the tear break-up time and a closer look at the tear break-up time of Hong Kong Chinese. *Optom Vis Sci* 1993;70(1):30-38.
- 21 McMonnies CW, Ho A. Patient history in screening for dry eye conditions. *J Am Optom Assoc* 1987;58(4):296-301.
- 22 Mengher LS, Bron AJ, Tonge SR, Gilbert DJ. A non-invasive instrument for clinical assessment of the pre-corneal tear film stability. *Curr Eye Res* 1985;4(1):1-7.
- 23 Efron N, Morgan PB, Katsara SS. Validation of grading scales for contact lens complications. *Ophthalmic Physiol Opt* 2001;21(1):17-29.
- 24 Itokawa T, Okajima Y, Suzuki T, Kakisu K, Iwashita H, Murakami Y, Hori Y. Association between ocular surface temperature and tear film stability in soft contact lens wearers. *Invest Ophthalmol Vis Sci* 2018;59(2):771-775.
- 25 Abusharha AA, Pearce EI, Fagehi R. Effect of ambient temperature on the human tear film. *Eye Contact Lens* 2016;42(5):308-312.
- 26 Buckmaster F, Pearce EI. Effects of humidity on tests of tear production. *Cornea* 2016;35(6):754-758.
- 27 Dursch TJ, Li W, Taraz B, Lin MC, Radke CJ. Tear-film evaporation rate from simultaneous ocular-surface temperature and tear-breakup area. *Optom Vis Sci* 2018;95(1):5-12.
- 28 Shen M, Wang J, Tao A, Chen Q, Lin S, Qu J, Lu F. Diurnal variation of upper and lower tear menisci. *Am J Ophthalmol* 2008;145(5):801-806.
- 29 Patel S, Bevan R, Farrell JC. Diurnal variation in precorneal tear film stability. *Am J Optom Physiol Opt* 1988;65(3):151-154.
- 30 Srinivasan S, Chan C, Jones L. Apparent time-dependent differences in inferior tear meniscus height in human subjects with mild dry eye symptoms. *Clin Exp Optom* 2007;90(5):345-350.
- 31 Tomlinson A, Cedarstaff TH. Diurnal variation in human tear evaporation. *J Br Contact Lens Assoc* 1992;15(2):77-79.
- 32 Packiriswamy V, Kumar P, Bashour M. Anthropometric and anthroposcopic analysis of periorbital features in Malaysian population: an inter-racial study. *Facial Plast Surg* 2018;34(4):400-406.
- 33 Li Q, Zhang X, Li K, Quan Y, Cai X, Xu S, Zhu F, Lu R. Normative anthropometric analysis and aesthetic indication of the ocular region for young Chinese adults. *Graefes Arch Clin Exp Ophthalmol* 2016;254(1):189-197.
- 34 Vasanthakumar P, Kumar P, Rao M. Anthropometric analysis of palpebral fissure dimensions and its position in South Indian ethnic adults. *Oman Med J* 2013;28(1):26-32.
- 35 The epidemiology of dry eye disease: report of the Epidemiology Subcommittee of the International Dry Eye WorkShop (2007). *Ocul Surf* 2007;5(2):93-107.
- 36 Schaumburg DA, Nichols JJ, Papas EB, Tong L, Uchino M, Nichols KK. The international workshop on meibomian gland dysfunction: report of the subcommittee on the epidemiology of, and associated risk factors for, MGD. *Invest Ophthalmol Vis Sci* 2011;52(4):1994-2005.
- 37 Truong S, Cole N, Stapleton F, Golebiowski B. Sex hormones and the dry eye. *Clin Exp Optom* 2014;97(4):324-336.
- 38 Craig JP, Tomlinson A. Importance of the lipid layer in human tear film stability and evaporation. *Optom Vis Sci* 1997;74(1):8-13.
- 39 Himebaugh NL, Begley CG, Bradley A, Wilkinson JA. Blinking and tear break-up during four visual tasks. *Optom Vis Sci* 2009;86(2):E106-E114.
- 40 Fagehi R, Ghazali H, Alrabiah S, Abusharha A, Alanazi S, Alsaqr A, Masmali A. Ocular dryness assessment in Saudi employees working indoors and outdoors. *Clin Optom (Auckl)* 2018;10:51-56.
- 41 Wolkoff P. Ocular discomfort by environmental and personal risk factors altering the precorneal tear film. *Toxicol Lett* 2010;199(3):203-212.