

Evaluation of intraocular penetration of levofloxacin by high performance liquid chromatography

Mahsa Hasanzadeh¹, Vafa Samarei², Amir Heydari^{3,4}

¹Department of Chemistry, Faculty of Science, Urmia University, Urmia 5756151818, Iran

²Department of Ophthalmology, Urmia University of Medical Sciences, Urmia 5715799313, Iran

³Center of Cellular and Molecular Research, Urmia University of Medical Science, Urmia 5715799313, Iran

⁴Department of Pharmacology, Faculty of Pharmacy, Urmia University of Medical Science, Urmia 5714783734, Iran

Correspondence to: Amir Heydari. Department of Pharmacology, Faculty of Pharmacy, Urmia University of Medical Science, Urmia, Iran. tmf_fmt3000@yahoo.com

Received: 2019-06-09 Accepted: 2020-02-29

Abstract

• **AIM:** To evaluate the levofloxacin eye drop into human eye penetration, levofloxacin eye drop concentrations in human ocular aqueous of 33 patients undergoing cataract surgery were measured by high performance liquid chromatography (HPLC).

• **METHODS:** Totally 33 volunteer patients who scheduled for phacoemulsification surgery received one drop of levofloxacin every 6h for 3d before and on the day of surgery, administration of drug was stopped 1h before surgery. Levofloxacin concentration in aqueous humor was measured by HPLC method with fluorescence detector.

• **RESULTS:** A simple, effective and sensitive HPLC method for determination of levofloxacin in human ocular aqueous was validated. Linearity was shown for levofloxacin concentration over a wide range of 1.95×10^{-3} -1.50 µg/mL. The mean aqueous level of levofloxacin was 0.3399 ± 0.03405 µg/mL.

• **CONCLUSION:** Results from the present study demonstrate that topical administration of levofloxacin 0.5% before cataract surgery with routine dose (one drop every 6h) unable to reach MIC90 for most common microorganism causing acute bacterial endophthalmitis.

• **KEYWORDS:** aqueous humour concentration; high performance liquid chromatography; levofloxacin; phacoemulsification

DOI:10.18240/ier.2021.03.07

Citation: Hasanzadeh M, Samarei V, Heydari A. Evaluation of intraocular penetration of levofloxacin by high performance liquid chromatography. *Int Eye Res* 2021;2(3):155-158

INTRODUCTION

Endophthalmitis is a rare but serious complication of cataract surgery, producing significant ocular morbidity and frequently irreversible visual loss. Most common causes of post-cataract endophthalmitis are coagulase-negative staphylococci (70%), *Staphylococcus aureus* (10%), streptococci (9%), other gram-positive cocci, including enterococci and mixed bacteria (5%), and gram-negative bacilli^[1-2]. Topical antibiotics preoperatively and postoperatively have been used to prevent this complication. Topical antibiotics might reduce ocular surface bacterial flora, but adequate penetration of antibiotics into anterior chamber (AC) is essential to reduce endophthalmitis. Fluoroquinolones are known for their broad spectrum of antibacterial activity against both gram-positive and gram-negative bacteria^[3]. Levofloxacin is a levo-isomer of the D, L-racemate ofloxacin compared with older-generation fluoroquinolones (ciprofloxacin, ofloxacin) it has broader spectrum and more activity against gram-positive and gram-negative bacteria. Levofloxacin has moderate bactericidal effect against anaerobes^[4-7]. Levofloxacin was determined with different methods like HPLC^[8-9], electrochemical^[10-12] and spectrophotometric method^[13]. The current study was designed to evaluate concentration of levofloxacin 0.5% in the aqueous humor after topical drop instillation in patients undergoing cataract surgery.

SUBJECTS AND METHODS

Ethical Approval The study was approved by an internal review board and informed consent was obtained from each patient.

The study group consisted of 33 patients (15 female and 18 male) with mean age of 64.5 ± 9.49 years who underwent elective phacoemulsification with intraocular lens implantation. The inclusion criteria included any patients with 18 years of age or older with significant cataract. Exclusion criteria included any patient with a known sensitivity to fluoroquinolones or who received topical ophthalmic

medications, except lubricants, during the month before surgery. All patients received one drop of medicine every six hours for three days before cataract surgery and on the day of surgery. Last administration was one hour before starting surgery. The pupils were dilated with tropicamide (0.5%). After preparation and draping of the eye, 0.1-0.15 mL of aqueous fluid was aspirated by paracentesis using a 30-gauge needle on a tuberculin syringe.

Levofloxacin hemihydrate was purchased from Aurobindo Co. Acetonitrile, triethylamine, and methanol were obtained from Merck (Germany). Orthophosphoric acid was purchased from BDH (England). All calibration standard samples were prepared every day. The mobile phase was isocratic consisting of 1% triethylamine (pH=3)/acetonitrile (86/14, v/v). The pH of 1% triethylamine was adjusted to 3.0 using phosphoric acid^[14]. The flow rate of mobile phase was modulated on 1.0 mL/min. The run time was 10min per each injection. 20 μ L was the volume of each sample that was injected. Manual sample injections were carried out using a Rheodyne model 7725I injector. Chromatographic analyses were carried out on a Cecil Adept system Binary Gradient liquid chromatography (Cecil, England) equipped with a two adept CE 4100 dual pistons pump and a Ultrafluor chrom Tech fluorescence Detector (Model LC 305, USA). Chromatographic separation was performed on HI-5, C18-100A (10 cm $\times$ 4.6 mm id.) reversed-phase column (Hichrom, England). The wavelengths of fluorescence detector were set to 294 nm (excitation) and 446 nm (emission).

All samples were stored and labeled in Eppendorf vials below -20°C. Then, all the collected samples were carried to the Center of Cellular and Molecular Research laboratory at the Urmia University of Medical Sciences for analysis. The samples were thawed, mixed for one minute and centrifuged for 10min at 3 000 g and 20 μ L of the clear supernatant injected into the column of HPLC analysis to assay levofloxacin concentrations.

RESULTS

Concentrations of levofloxacin were successfully measured by high performance liquid chromatography (HPLC) method without any interfering peaks. Chromatogram for the assay of levofloxacin concentration was shown on Figure 1. As observed in Figure 2 according to calibration curve, there is a linear relationship between the concentration of levofloxacin and peak area in a wide range of 1.95×10^{-3} -1.50 μ g/mL. The equation of the calibration curve was as follows: $Y=284.62X+0.6742$ with a correlation coefficient (r^2) of 0.9997, where Y is the peak area and X is the concentration of levofloxacin (μ g/mL). Mean concentration was 0.3399 ± 0.03405 μ g/mL.

Table 1 shows the precision and accuracy of the method. The coefficient of variation percent (CV%) ranged from 2.37-1.91




Figure 1 Representing the chromatogram correlates to levofloxacin with excitation and emission wavelengths 24 and 446 sequence. Moreover, retention time 05:42.4min Showing levofloxacin with a concentration of 0.5 μ g/mL.




Figure 2 Calibration plot corresponds to concentration of levofloxacin.




Figure 3 Concentration of levofloxacin in aqueous humor after administration of levofloxacin 0.5%.

for intra-day experiments that were repeated three times on the same day. CV% for inter-day experiments that were repeated over a period of three consecutive days was calculated in the range of 0.89-2.93. The accuracy of the assay ranged from 94.37-102.82 for intra-day experiments and 94.40 to 102.82 for inter-day analysis. Therefore, the proposed method possesses satisfying precision and accuracy. Concentration of levofloxacin in human ocular aqueous was illustrated in Figure 3. No adverse reactions were reported after the use of levofloxacin drops.

Table 1 Intra-day and inter-day precision and accuracy data for evaluation of levofloxacin concentration by HPLC

Concentration (µg/mL)	Intra-day precision CV%	Intra-day accuracy %	Inter-day precision CV%	Inter-day accuracy %
0.125	2.27	94.37	2.93	94.40
0.25	2.37	102.08	0.89	102.08
0.5	1.91	102.82	1.33	102.82
Mean	2.14	100.50	1.72	99.77

DISCUSSION

Endophthalmitis typically presents as a moderate to severe infection 5 to 7 days after surgery. Less commonly a chronic, indolent form presents several months after surgery and is usually caused by *P. acnes*. Post-operative bacterial endophthalmitis is caused by gram-positive microbes in majority of isolates. Infection that caused by streptococci and Gram-negative bacteria such as *Pseudomonas aeruginosa* have a poor Visual prognosis^[1-2]. The level of levofloxacin is unable to reach minimum inhibitory concentration 90 (MIC90) for most common microorganism responsible for endophthalmitis (Table 2). Despite the use of the best available treatments, the visual prognosis of patients with intraocular infections remains guarded^[17-18]. Levofloxacin possesses a wide spectrum of bactericidal activity against both gram-positive and gram-negative bacteria, as well as a typical pathogens such as *Mycoplasma*, *Chlamydia* and *Legionella*. Concerning ocular penetration, levofloxacin appears to have several advantages over other fluoroquinolones. Levofloxacin is highly water soluble, which enables the aqueous solution to be as high as 2% at neutral pH (35.8 mg/mL) and for this reason high concentration aqueous solutions of levofloxacin commercially available (0.5%). Furthermore, because levofloxacin consists of only the active L-isomer of the D- and L-racemate ofloxacin, levofloxacin is roughly twice as biologically active as other antibiotics in the anterior chamber^[7,19-20].

In other studies results were higher than mean levofloxacin concentration in our study. These studies used pulsed delivery levofloxacin 0.5% before cataract surgery. (For example every 10 minutes for 5 doses^[21] or every 15min for 4 doses^[22] Mean levofloxacin concentration in pulsed delivery was much higher than our report. $1.135 \pm 0.589 \mu\text{g/mL}$ ^[21] and $(1.139.9 \pm 0.717 \mu\text{g/mL})$ respectively^[22]. Sundelin *et al*^[23] in 2009 reported the higher anterior chamber level of levofloxacin, because they were administered the medicine as pulsed methods.

However, the difference in the anterior chamber concentrations might be mainly attributed to the difference in the concentration of the administered eye drop solutions. The previous studies have showed that frequent dosing of the antibiotics on the day of surgery give the highest concentration in the anterior chamber^[23]. The frequent dosing of levofloxacin 0.5% eye drops immediately before or after surgery helps prevent the development of postoperative endophthalmitis.

Table 2 Representative MIC₉₀ Values for Staphylococcus and Streptococcus Strains reported for levofloxacin and Ciprofloxacin for several ocular pathogens

Organisms	MIC ₉₀ (µg/mL)		
	Levofloxacin	Ciprofloxacin	References
<i>S. aureus</i>	0.5	1.0	[15]
	1.0	1.0	[16]
<i>S. epidermidis</i>	0.5	1.0	[15]
	1.0	1.0	[16]
<i>S. pneumoniae</i>	1.0	2.0	[15]
	2.0	-	[16]
<i>S. pyogenes</i>	2.0	-	[16]
	1.0	1.0	[15]
<i>S. agalactiae</i>	1.0	1.0	[15]
	2.0	-	[16]

Koch *et al*^[22] showed that the concentration of the drug in the anterior chamber correlated negatively with the anterior chamber depth and volume, which must consider to explain the low concentration of levofloxacin in my studies. Some studies detected correlation between fluoroquinolone concentrations and the corneal diameters or corneal surface areas might have influenced drug uptake.

Kessel *et al*^[24] published a review article deals with the use of topical antibiotics in the prevention of endophthalmitis. Three days of topical antibiotic treatment reduces the number of positive conjunctival samples by approximately 50%. It means after topical antibiotic a significant number of bacteria remain in the ocular surface treatment reduces the number of positive conjunctival samples by approximately 50%^[24]. Topical antibiotic therapy was not found to lower the rate of endophthalmitis in the European Society of Cataract and Refractive Surgeons (ESCRS) study^[25]. In the Conclusions, the researcher reported that could not find any evidence that topical antibiotic treatment after cataract surgery lowers the risk of endophthalmitis. As there is no documented effect of topical antibiotic treatment and its use may be associated with concern for selection of resistant bacterial strains, we cannot recommend using it^[24].

Results from the present study demonstrate that topical administration of levofloxacin 0.5% before cataract surgery with routine dose (one drop every 6h) fail to reach MIC90 for most common microorganism causing acute bacterial endophthalmitis. Thus, further researches required before

prescription of these two drugs. Clinical application of routine dose of levofloxacin that more than 80%-90% of ophthalmologists use it, can't be trustworthy for prevention of discussed infections.

ACKNOWLEDGEMENTS

Conflicts of Interest: Hasanazadeh M, None; Samarei V, None; Heydari A, None.

Peer Review File Available at: http://ier.ijo.cn/gjykier/ch/reader/download_attache_file.aspx?seq_id=20210924103808001&flag=1&journal_id=gjykier&year_id=2021&issue=3

REFERENCES

- 1 Endophthalmitis Study Group, European Society of Cataract & Refractive Surgeons. Prophylaxis of postoperative endophthalmitis following cataract surgery: results of the ESCRS multicenter study and identification of risk factors. *J Cataract Refract Surg* 2007;33(6):978-988.
- 2 Durand ML. Endophthalmitis. *Clin Microbiol Infect* 2013;19(3):227-234.
- 3 Graves A, Henry M, O'Brien TP, Hwang DG, van Buskirk A, Trousdale MD. *In vitro* susceptibilities of bacterial ocular isolates to fluoroquinolones. *Cornea* 2001;20(3):301-305.
- 4 van Bambeke F, Michot JM, van Eldere J, Tulkens PM. Quinolones in 2005: an update. *Clin Microbiol Infect* 2005;11(4):256-280.
- 5 Sasaki K, Yukihiro M, Masamichi F, Masao O, Yuichi O. Intraocular penetration mode of five fluoroquinolone ophthalmic solutions evaluated by the newly proposed parameter of AQC_{max} . *J Eye* 1995;12(5):787-790.
- 6 Davis R, Bryson HM. Levofloxacin. A review of its antibacterial activity, pharmacokinetics and therapeutic efficacy. *Drugs* 1994;47(4):677-700.
- 7 Yamada M, Mochizuki H, Yamada K, Kawai M, Mashima Y. Aqueous humor levels of topically applied levofloxacin, norfloxacin, and lomefloxacin in the same human eyes. *J Cataract Refract Surg* 2003;29(9):1771-1775.
- 8 Szerkus O, Jacyna J, Gibas A, Sieczkowski M, Siluk D, Matuszewski M, Kaliszan R, Markuszewski MJ. Robust HPLC-MS/MS method for levofloxacin and ciprofloxacin determination in human prostate tissue. *J Pharm Biomed Anal* 2017;132:173-183.
- 9 van Toi P, Pouplin T, Tho NDK, Phuong PN, Chau TTH, Thuong Thuong NT, Heemskerck D, Hien TT, Thwaites GE. High-performance liquid chromatography with time-programmed fluorescence detection for the quantification of Levofloxacin in human plasma and cerebrospinal fluid in adults with tuberculous meningitis. *J Chromatogr B Analyt Technol Biomed Life Sci* 2017;1061-1062:256-262.
- 10 Wong A, Santos AM, Fatibello-Filho O. Simultaneous determination of paracetamol and levofloxacin using a glassy carbon electrode modified with carbon black, silver nanoparticles and PEDOT: PSS film. *Sens Actuat B: Chem* 2018;255:2264-2273.
- 11 Rkik M, Brahim MB, Samet Y. Electrochemical determination of levofloxacin antibiotic in biological samples using boron doped diamond electrode. *J Electroanal Chem* 2017;794:175-181.
- 12 Han L, Zhao YF, Chang C, Li F. A novel electrochemical sensor based on poly(p-aminobenzene sulfonic acid)-reduced graphene oxide composite film for the sensitive and selective detection of levofloxacin in human urine. *J Electroanal Chem* 2018;817:141-148.
- 13 El-Brashy AM, Metwally MS, El-Sepai FA. Spectrophotometric determination of some fluoroquinolone antibacterials by ion-pair complex formation with cobalt (II) tetrathiocyanate. *J Chin Chem Soc* 2013;52(1):77-84.
- 14 Watabe S, Yokoyama Y, Nakazawa K, Shinozaki K, Hiraoka R, Takeshita K, Suzuki Y. Simultaneous measurement of pazufloxacin, ciprofloxacin, and levofloxacin in human serum by high-performance liquid chromatography with fluorescence detection. *J Chromatogr B Analyt Technol Biomed Life Sci* 2010;878(19):1555-1561.
- 15 Pickerill KE, Paladino JA, Schentag JJ. Comparison of the fluoroquinolones based on pharmacokinetic and pharmacodynamic parameters. *Pharmacotherapy* 2000;20(4):417-428.
- 16 Patel JB, Weinstein MP, Eliopoulos GM, Jenkins SG, Lewis JS, Limbago B, Mathers AJ, Mazzulli T, Patel R, Richter SS, Satlin M, Swenson JM, Traczewski MM, Turnidge JD, Zimmer BL. Performance standards for antimicrobial susceptibility testing. 27th Edition. *USA:CLSI* 2017.
- 17 Benz MS, Scott IU, Flynn HW, Unonius N Jr, Miller D. Endophthalmitis isolates and antibiotic sensitivities: a 6-year review of culture-proven cases. *Am J Ophthalmol* 2004;137(1):38-42.
- 18 Callegan MC, Engelbert M, Parke DW 2nd, Jett BD, Gilmore MS. Bacterial endophthalmitis: epidemiology, therapeutics, and bacterium-host interactions. *Clin Microbiol Rev* 2002;15(1):111-124.
- 19 North DS, Fish DN, Redington JJ. Levofloxacin, a second-generation fluoroquinolone. *Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy* 1998;18(5):915-935.
- 20 Colin J, Simonpoli S, Geldsetzer K, Ropo A. Corneal penetration of levofloxacin into the human aqueous humour: a comparison with ciprofloxacin. *Acta Ophthalmol Scand* 2003;81(6):611-613.
- 21 Bucci FA. An *in vivo* study comparing the ocular absorption of levofloxacin and ciprofloxacin prior to phacoemulsification. *Am J Ophthalmol* 2004;137(2):308-312.
- 22 Koch HR, Kulus SC, Roessler M, Ropo A, Geldsetzer K. Corneal penetration of fluoroquinolones: aqueous humor concentrations after topical application of levofloxacin 0.5% and ofloxacin 0.3% eyedrops. *J Cataract Refract Surg* 2005;31(7):1377-1385.
- 23 Sundelin K, Seal D, Gardner S, Ropo A, Geldsetzer K, Lökkila J, Stenevi U. Increased anterior chamber penetration of topical levofloxacin 0.5% after pulsed dosing in cataract patients. *Acta Ophthalmol* 2009;87(2):160-165.
- 24 Kessel L, Flesner P, Andresen J, Erngaard D, Tendal B, Hjortdal J. Antibiotic prevention of postcataract endophthalmitis: a systematic review and meta-analysis. *Acta Ophthalmol* 2015;93(4):303-317.
- 25 Råen M, Sandvik GF, Drolsum L. Endophthalmitis following cataract surgery: the role of prophylactic postoperative chloramphenicol eye drops. *Acta Ophthalmol* 2013;91(2):118-122.